

OPERATION OF THE PRIOR INFORMED  
CONSENT PROCEDURE FOR BANNED  
OR SEVERELY RESTRICTED CHEMICALS  
IN INTERNATIONAL TRADE

# DECISION GUIDANCE DOCUMENTS

**Pentachlorophenol and its salts and esters**

JOINT FAO/UNEP PROGRAMME  
FOR THE OPERATION OF  
PRIOR INFORMED CONSENT



United Nations Environment Programme



Food and Agriculture Organization  
of the United Nations

**OPERATION OF THE PRIOR INFORMED CONSENT PROCEDURE FOR BANNED  
OR SEVERELY RESTRICTED CHEMICALS IN INTERNATIONAL TRADE**

DECISION GUIDANCE  
DOCUMENTS

Pentachlorophenol and its salts and esters

JOINT FAO/UNEP PROGRAMME FOR THE OPERATION OF  
PRIOR INFORMED CONSENT

Food and Agriculture Organization of the United Nations  
United Nations Environment Programme  
Rome - Geneva 1991; amended 1996

## **DISCLAIMER**

The inclusion of these chemicals in the Prior Informed Consent Procedure is based on reports of control action submitted to the United Nations Environment Programme (UNEP) by participating countries, and which are presently listed in the UNEP-International Register of Potentially Toxic Chemicals (IRPTC) database on Prior Informed Consent. While recognizing that these reports from countries are subject to confirmation, the FAO/UNEP Joint Working Group of Experts on Prior Informed Consent has recommended that these chemicals be included in the Procedure. The status of these chemicals will be reconsidered on the basis of such new notifications as maybe made by participating countries from time to time.

The use of trade names in this document is primarily intended to facilitate the correct identification of the chemical. It is not intended to imply approval or disapproval of any particular company. As it is not possible to include all trade names presently in use, only a number of commonly used and published trade names have been included here.

This document is intended to serve as a guide and to assist authorities in making a sound decision on whether to continue to import, or to prohibit import, of these chemicals because of health or environmental reasons. While the information provided is believed to be accurate according to data available at the time of preparation of this Decision Guidance Document, FAO and UNEP disclaim any responsibility for omissions or any consequences that may flow therefrom. Neither FAO or UNEP, nor any member of the FAO/UNEP Joint Group of Experts shall be liable for any injury, loss, damage or prejudice of any kind that may be suffered as a result of importing or prohibiting the import of these chemicals.

The designations employed and the presentation of material in this publication do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations or the United Nations Environment Programme concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

## ABBREVIATIONS WHICH MAY BE USED IN THIS DOCUMENT

(N.B. : chemical elements and pesticides are not included in this list)

ADI	acceptable daily intake
ai	active ingredient
b.p.	boiling point
bw	bodyweight
°C	degree Celsius (centigrade)
CCPR	Codex Committee on Pesticide Residues
DNA	Designated National Authority
EC	emulsion concentrate
EEC	European Economic Community
EPA	U.S. Environmental Protection Agency
ERL	extraneous residue limit
FAO	Food and Agriculture Organization of the United Nations
g	gram
µg	microgram
GAP	good agricultural practice
GL	guideline level
ha	hectare
IARC	International Agency for Research on Cancer
i.m.	intramuscular
i.p.	intraperitoneal
IPCS	International Programme on Chemical Safety
IRPTC	International Register of Potentially Toxic Chemicals
JMPR	Joint FAO/WHO Meeting on Pesticide Residues (Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues)
k	kilo- (x 10 <sup>3</sup> )
kg	kilogram
l	litre
LC <sub>50</sub>	lethal concentration, 50%
LD <sub>50</sub>	lethal dose, median

m	metre
mg	milligram
ml	millilitre
m.p.	melting point
MRL	Maximum Residue Limit.
MTD	maximum tolerated dose
ng	nanogram
NOEL	no-observed-effect level
NOAEL	no-observed-adverse-effect level
NS	Not Stated
OP	organophosphorus pesticide
PHI	pre-harvest interval
ppb	parts per billion
ppm	parts per million (Used only in reference to the concentration of a pesticide in an experimental diet. In all other contexts the terms mg/kg or mg/l are used).
ppt	parts per trillion
sp gr	specific gravity
STEL	Short Term Exposure Limit
TADI	Temporary Acceptable Daily Intake
TLV	Threshold Limit Value
TMDI	theoretical maximum daily intake
TMRL	Temporary Maximum Residue Limit
TWA	Time Weighted Average
UNEP	United Nations Environment Programme
WHO	World Health Organization
WP	wettable powder
wt	weight
<	less than
<<	much less than
≤	less than or equal to
>	greater than
≥	greater than or equal to

## Prior Informed Consent Decision Guidance Document

# Pentachlorophenol

## 1 Identification

1.1	<b>Common Name</b>	Pentachlorophenol and its salts and esters
	<b>Other names/synonyms</b>	PCP, penta, penchlorol, chlorophen
1.2	<b>Chemical Type</b>	Chloronitrophenol derivative
1.3	<b>Use</b>	Wood use: algaecide, fungicide, insecticide (wood preservative) Non-wood use: general disinfectant, herbicide, insecticide (termiticide), molluscicide, anti-fouling paint
1.4	<b>Chemical Name</b>	Pentachlorophenol
1.5	<b>CAS No.</b>	87-86-5
1.6	<b>Trade Names</b>	Block Penta; Chem-Tol; Cryptogil oil; Dowcide 7/EC-7/G; Dowicide G; Dirottox; EP 30, Fungifen; GLAZD Penta; Grundier Arbezol; Lautor A, Lauxtol; Lauxtrol A; Lipoprem; Pentchloral; Pentacon; Penta C 30; Penta-Kil; Penta Plus 40; Penta Pres 1-10; Penta WR1-5 Penwar; Peratox; Permacide; Permagard; Persasan; Permattox; Permite; Santobrite; Santophen; Santophen 20; Sauttox, Sinituho; Term-i-Trol; Thompson's Wood Fix
1.7	<b>Mode of action as Pesticide</b>	It is widely believed that pentachlorophenol affects uncoupling mitochondrial oxidative phosphorylation, thereby causing accelerated aerobic metabolism and increasing heat production. It causes loss of membrane electrical resistance
1.8	<b>Formulation Types</b>	Wettable powder (50%), prills (or granular) (85-90%), blockform 90+%, oil-miscible liquid (5%), emulsifiable concentrate (40%). PCP is available as a sodium salt. The compound may be used alone or in combination with other agents, such as sodium fluoride, dichromate salts, sodium arsenate or arsenious oxide (Farm Chemicals Handbook, 1991).
1.9	<b>Basic Manufacturers</b>	Vulcan Materials Co., (Chemicals Div.), Birmingham, Alabama, USA. In the 1980s produced by Ameco, Canada; National Product Co., China; Chapman Chemicals; KMG Mernuth (Mexico); Pola Quimia SA de CB, Mexico City, Mexico; Preservation Products, Matamoros, Mexico; Melchemie, Holland; Rhône-Poulenc, Lyons, France

## 2 Summary of Control Actions

## 2.1 **General**

Control actions to ban or severely restrict pentachlorophenol have been reported by eight countries and the European Union. Six have banned the pesticide; two and the EU have severely restricted it.

*The control actions reported by governments to IRPTC/UNEP are listed in Annex 1.*

## 2.2 **Reasons for the Control Actions**

Pentachlorophenol has been subject to control actions primarily because of high toxicity to human and animals. Developmental and reproductive effects and liver and kidney pathology were noted in animal studies. However, the compound is also highly toxic to aquatic organisms. Pentachlorophenol contains several highly toxic dioxins which have shown carcinogenic effects in experimental animals.

## 2.3 **Bans and restrictions**

Pentachlorophenol continues to be used extensively. However, depending upon the reporting country, the number of banned uses ranges from all uses to few uses. Most reporting countries banned residential indoor uses. Austria, India, Indonesia, New Zealand, Sweden and Switzerland have reported a total ban.

*Refer to Annex 1 for details.*

## 2.4 **Uses Reported to be Continued in Effect**

Belize and China have retained uses for wood preservation purposes by licensed applicators or under specific conditions.

## 2.5 **Alternatives**

A number of chemical alternatives to PCP for wood preservation have been developed. While many of these, such as TCMTB, are as toxic as or more toxic than purified PCP, they do not contain the persistent toxic contaminants that characterize formulated PCP products. Other, less toxic PCP alternatives now in use include copper naphthanate and zinc naphthanate. PCP has been replaced by other chemical agents for virtually all of its former agricultural uses. Specific alternatives were suggested by India. Many less toxic alternatives are registered in Indonesia and Austria. For details, see Annex 2.

*It is important to remember that the effectiveness of any alternative pesticide needs to be established under conditions of use in specific crops and countries.*

## 2.6 **Contacts for Further Information**

*FAO/UNEP Joint Data Base, IRPTC, Geneva; Designated National Authorities (DNAs) in countries taking control actions may be a source of information on alternatives (Annex 3).*

# **3 Summary of Further Information on**

## Pentachlorophenol

### 3.1 Chemical and Physical Properties

Pure pentachlorophenol forms colourless monoclinic crystals. The technical grade is a dark-grey to brown pellet/powder. Melting point is 191°C (anhydrous), 174°C (monohydrous). Boiling point (decomposition) 309-310°C. The vapour pressure at 20°C is 2 mPa. Solubility (mg/l) water: 20 mg/l at 30 °C. It is freely soluble in organic solvents, and alcohol, soluble in benzene, ethanol and diethyl ether; slightly soluble in cold petroleum ether; carbon tetrachloride and paraffins; inflammable. Commercial PCP contains very toxic impurities. Those of greatest concern are hexachlorodibenzo-p-dioxin (HxCDD), 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), the chlorinated dibenzofurans and hexachlorobenzene (HCB) (Royal Society of Chemistry, 1991; Pesticide Manual, 1991).

### 3.2 Toxicological Characteristics

#### 3.2.1 Classification

- WHO** Class Ib ( highly hazardous :WHO, 1994)  
**EU** Carcinogen (Cat.3): highly toxic; irritant  
**IARC** No evaluation of carcinogenicity because data are inadequate

#### 3.2.2 General

PCP is highly toxic to mammals and birds. It is the most acutely toxic of the chlorophenols tested. Some of the acute effects of exposure to commercial PCP are attributable to microcontaminants present in the technical preparation (IPCS, 1987).

**Uptake** PCP is readily absorbed when applied to the skin

#### 3.2.3 Acute Toxicity

- Oral** LD<sub>50</sub>: 27-205 mg/kg (tests with a variety of species) (IPCS, 1987)  
**Inhalation** Rabbit (35.2%) LD<sub>50</sub> 201 mg/kg  
**Dermal** LD<sub>50</sub>: 80-350 mg/kg (tests with a variety of species) (IPCS, 1987)  
**Irritation** Irritating to skin, eyes and mucous membranes (US PHS, 1989; IPCS, 1987)

#### 3.2.4 Short-term Toxicity

**Reproduction** Numerous studies have described the developmental effects of pentachlorophenol and its dioxin and hexachlorobenzene contaminants. PCP is foetotoxic and teratogenic when administered during early gestation (Eisler, 1989). There is a general agreement that PCP is a foetotoxic agent; however, it does not appear to be teratogenic (NRCC, 1982). It was concluded that the NOEL for teratogenicity, foetotoxicity and embryotoxicity in rats was 10 mg/kg.

**NOEL** The NOEL was determined to be 1.25 mg/kg/day (25 ppm, LDT) subchronic inhalation (rat) study.

#### 3.2.5 Chronic Toxicity

**Carcinogenicity** PCP was tested for carcinogenicity by administration in the diet of two strains of mice and in one experiment in rats (IARC, 1979). The

data available on the carcinogenic properties of PCP were reviewed and it was concluded that no carcinogenic effects were evident in either species. The IARC therefore concluded that there is inadequate evidence for carcinogenicity to animals (IARC, 1987). A review of data by the Carcinogenic Assessment Group of the US EPA concluded that PCP was negative with respect to oncogenic effects (Williams, 1982).

The same studies in mice (Innes *et al.*, 1969;) did not indicate significant increases in tumour incidence.

**Mutagenicity** PCP probably does not cause mutations. The data available are insufficient to fully assess the mutagenicity of PCP (Williams, 1982).

**Others** PCP has proved to be immunotoxic for mice, rats, chicken and cattle. Neurotoxic effects have also been reported (IPCS, 1987).

**3.2.6 Epidemiological Data** Most of the available information regarding effects of pentachlorophenol in humans comes from cases of acute over-exposure following the home use of PCP in wood preservation and herbicides and occupational exposure in agriculture and the wood treatment industry. The few available industrial surveys and epidemiological studies are limited in their usefulness because of small sample size, short follow-up periods and brief exposure periods. Nevertheless, these studies suggest that PCP can adversely affect the liver, kidney, skin, blood, lungs and central nervous system.

### 3.3 Environmental Characteristics

**3.3.2 Effects** PCP is highly toxic to aquatic organisms. Invertebrates and fish are adversely affected by concentrations of PCP below 1 mg/l; algae are very sensitive to PCP.

**Fish** PCP LC<sub>50</sub>: Bluegill 23-92.5 µg/L., Rainbow trout 48-68.7 µg/l

**Aquatic Invertebrates** Eisler (1989) reported a 48-hour median lethal concentration (LC<sub>50</sub>) of 260 µg/l for the clawed toad

**Birds** Avian toxicity may be somewhat less sensitive to PCP than mammalian: LD<sub>50</sub>s 380 mg/kg bw for mallard duck and 504 mg/kg bw for ring-necked pheasant

### 3.4 Exposure

- 3.4.1 Food/drinking water** Low use in foods. However, in countries where large amounts of PCP have been used, contamination of foods and feed commodities is probably from environmental movement. Therefore, consumption of contaminated foods represents the primary route for most people (UN ECE, 1994). Overall estimates of PCP intake from all foods, based on total diet samples in the USA and Germany, range up to approximately 6 µg/person per day (IPCS, 1989).
- 3.4.2 Occupational/Use** The extensive use of PCP to treat wood, and to a lesser extent use in homes and gardens, together with its physical and fate characteristics indicate that there is likely to be widespread human exposure occurring partially through skin contact, but mainly through inhalation, which is the most dangerous route of exposure to PCP. This is confirmed by many reports of its occurrence in the general environment and its presence in body-fluids, both in the general population and in exposed workers (Fielders *et al.*, 1982). Airborne levels of PCP production and wood-preservation facilities have ranged from several mg/m<sup>3</sup> to more than 300 mg/m<sup>3</sup> in some work areas. Under these circumstances, the ADI may be significantly exceeded.  
Domestic use, such as indoor application of wood preservatives and paints based on PCP or PCP-treated wood or indoor wood panels or boards, leads to high concentrations in the indoor atmosphere, and the ADI may be significantly exceeded.
- 3.4.3 Environment** The relatively high volatility and mobility of PCP and the water solubility of its ionized form have led to the widespread contamination of all environmental sectors, and a long-range dissemination of this compound. PCP will leach from treated wood, volatilise from treated surfaces and may get into waterways, adversely affecting fish.
- 3.4.4 Accidental Poisoning** IARC (1979) reports one case of fatal aplastic anaemia from exposure to pentachlorophenol and tetrachlorophenol and the death of nine sawmill workers exposed to treated wood. Two deaths were reported among 20 infants intoxicated in a hospital due to the misuse of a laundry product containing 22.9% sodium penta-chlorophenol, 4% 3,4,4-trichlorocarbanilide and sodium salts of other chlorophenols and inert ingredients. Extended periods of exposure to PCP have resulted in persistent chloracne and disorders of the nervous system and liver. There are first aid treatments available for ingestion, inhalation and contamination of the eye and skin. Notes to physicians are also available.

### 3.5 Measures to Reduce Exposures

Exposure may be reduced by providing protective clothing. Where dermal contact is expected wear gloves. Spray applicators should wear protective clothing (e.g. overalls, jacket, gloves and boots) which is impervious to wood treatment formulations, plus a respirator, head covering and goggles when spraying. Keep clothes clean or dispose of clothing. Do not eat, drink or smoke when contaminated from applying PCP. Automated processes and the use of closed systems, where applied, have greatly reduced worker-exposure.

### 3.6 Packaging and Labelling

The product should be labelled explicitly. Handling instructions are required. For further advice refer to the FAO Revised Guidelines on Good Labelling Practice for Pesticides (1995).

### 3.7 Waste Disposal Methods (WHO/IPCS, 1990)

The disposal of technical PCP and associated waste should preferably involve controlled high-temperature combustion with effluent gas scrubbing to prevent release of hydrogen chloride gas.

### 3.8 Maximum Residue Limits (mg/kg)

**Codex/JMPR ADI:** The Codex/JMPR has no established ADI for Pentachlorophenol

**Codex MRLs:** The Codex/JMPR has no established MRLs for Pentachlorophenol

**Belgium:** Maximum permissible levels in mg/kg: mushrooms 0.05; other 0\* (0.01).

\* ISO designation

**Germany** (Maximum levels in mg/kg) All foods of plant origin 0.01.

**Israel** (Maximum residue limits in mg/kg.) Mushrooms and other foods 0.05.

**Netherlands** (Maximum residue limits in mg/kg) Mushroom 0.05; other 0\* (0.01\*).

Note: (\*) residues shall be absent, while the highest concentration at which this requirement is still deemed to have been met is indicated in parentheses

**Switzerland** (Limit value in mg/kg) Milk 0.05.

**Yugoslavia** (Tolerance level in ppm) Unnamed commodities 0.01.

## 4 Major References

**CIRAD (1990).** Agricultural Requisites Scheme for Asia and the Pacific & International Co-operation Centre of Agricultural Research for Development. The ARSAP/CIRAD regional agro-pesticide index Asia

**Eisler, R. (1989).** Pentachlorophenol hazards to fish, wild and invertebrates: A synoptic review. Patuxent Wildlife Research Centre. Laurel, MD, U.S. Fish and Wildlife Service. 72 pp

**Farm Chemicals Handbook '90 (1990).** Meister Publishing, Willoughby, OH, USA

**Fielders et al. (1982).** Pentachlorophenol. *In:* Toxicity review, London, Health & Safety Executive, Vol., 20 pp

**FAO (1995).** Revised guidelines on good labelling practices for pesticides. Food and Agriculture Organization, Rome

**FAO (1996).** Technical guidelines on disposal of bulk quantities of obsolete pesticides in developing countries. Food and Agriculture Organization, Rome

**Health Welfare Canada (1990).** National pesticide residue limits in food. Ottawa, Ontario, Canada

- Haley, T. (1977).** Human poisoning with pentachlorophenol and its treatment. *Ecotoxicology and Environmental Safety* Vol. 1
- Innes, J.R.M. et al. (1969).** Bioassay of Pesticides and Industrial Chemicals for Tumorigenicity in mice: A preliminary Note. *J. Natl. Cancer Inst.* 42: 1101-1114
- IARC (1979).** IARC monographs on the evaluation of the carcinogenic risk of humans, Volume 20. International Agency for Research on Cancer, Lyons, France
- IARC (1987).** IARC monographs on the evaluation of the carcinogenic risk of humans, Supplement 7. International Agency for Research on Cancer, Lyons, France
- IPCS (1987).** Environmental Health Criteria No. 71. International Programme on Chemical Safety, WHO, Geneva
- IPCS (1989).** IPCS Health and Safety Guide Series No. 19. International Programme on Chemical Safety, WHO, Geneva
- NRCC (1982).** Chlorinated phenols: criteria for environmental quality. National Research Council of Canada, Associate Committee on Scientific Criteria for Environmental Quality, Ottawa, Ontario, No. 18578) 191 pp
- Royal Society of Chemistry (1991).** The Agrochemicals Handbook (3rd ed.). Cambridge, United Kingdom
- U.S. Environmental Protection Agency (1987).** Determination and intent to cancel and deny applications for registrations of pesticide products containing pentachlorophenol. *Federal Register*, Vol. 52, No. 13, Government Printing Office, Washington, DC (January 21, 1987)
- U.S. Environmental Protection Agency (1988).** Pentachlorophenol products; amendment of notice of intent to cancel registrations of products for non-wood biocide uses. *Federal Register*, Vol. 53, No. 36, Government Printing Office, Washington, DC (February 24, 1988)
- U.S. Public Health Service/Environmental Protection Agency (1989).** The toxicological profile for pentachlorophenol. USPHS & USEPA, Washington (Dec. 1989)
- Williams, P.L. (1982).** Pentachlorophenol: an assessment of the occupational hazard. *Am. Ind. Hyg. Assoc. J.*, 43:799-810
- WHO (1996).** The World Health Organization Recommended Classification of Pesticides by Hazard and Guidelines to Classification 1996-1997. World Health Organization, Geneva, WHO/PCS/96.3
- Worthing, C.R. and R.J. Hance (Eds.) (1994).** The Pesticide Manual: A World Compendium. (10th ed.). British Crop Protection Council, Surrey, United Kingdom

## ANNEX 1

**Summary of Control Actions and Remaining Uses as Reported by Countries****Actions taken and year effective****Bans**

<b>Austria</b>	
<b>Control Action</b> <b>Effective</b> <b>Uses still allowed</b> <b>Reasons for control action</b>	Pentachlorophenol and its salts are banned 1991 Use is still allowed for scientific and analytical purposes Highly toxic impurities in commercial products (chlorinated dibenz o-p-dioxin: CDDs) and chlorinated dibenzofurans (CDFs) and formation of highly toxic compounds on combustion. CDDs and CDFs have shown carcinogenic effects in experimental animals.
<b>India</b>	
<b>Control Action</b> <b>Effective</b> <b>Uses still allowed</b> <b>Reasons for control action</b>	Banned 1991  Owing to high toxicity to man, animals and aquatic organisms and presence of toxic impurities in commercial products
<b>Indonesia</b>	
<b>Control Action</b> <b>Effective</b> <b>Uses still allowed</b> <b>Reasons for control action</b>	Prohibited for all uses 1980 No remaining uses allowed Extremely toxic; may be fatal if swallowed or absorbed through skin; causes skin irritation; vapours will cause injury
<b>New Zealand</b>	
<b>Control Action</b>      <b>Effective</b> <b>Uses still allowed</b> <b>Reasons for control action</b>	1) All uses and products are banned 2) Agreement in principle to permit re-introduction in closed timber treatment systems at approved sites with specific conditions on disposal of waste. These conditions have not been met and therefore no products registered, no use permitted and no imports allowed 1991 No remaining uses allowed
<b>Sweden</b>	

<b>Control Action</b> <b>Effective</b> <b>Uses still allowed</b> <b>Reasons for control action</b>	Pentachlorophenol and other chlorophenols are banned 1978 No remaining uses allowed. Because of highly toxic impurities in commercial products
---	---

<b>Switzerland</b>	
<b>Control Action</b>  <b>Effective</b> <b>Uses still allowed</b> <b>Reasons for control action</b>	Totally banned chemical: manufacture, supply, import and use of the substance and of products which contain the substance are prohibited. (Applies to pentachlorophenol, its salts and pentachlorophenoxy compounds) 1988 No remaining uses allowed Bioaccumulation, highly toxic impurities, formation of highly toxic substances on thermolysis

## Severely Restricted

<b>EU/EEA<sup>1</sup></b>	
<b>Control Action</b>  <b>Effective</b> <b>Uses still allowed</b>         <b>Reasons for control action</b>	Pentachlorophenol, its salts and esters shall not be used in a concentration equal to or greater than 0.1% by mass in substances or preparations placed on the market for use by the general public. 1992 Does not apply to substances and preparations for use in industrial installations with emission and/or discharge of PCP greater than those prescribed by existing legislation; and only (a) in the treatment of wood neither intended for use inside buildings, nor for manufacture of containers intended for growing purposes, nor manufacture of packaging for products intended for human and/or animal consumption; (b) in the impregnation of fibres and heavy-duty textiles not intended for clothing or for decorative furnishing; (c) as a synthesizing and/or processing agent in industrial processes; (d) by way of special exception for remedial treatment of timber and masonry infected by dry rot fungus ( <i>Serpula lacrymans</i> ). PCP, its salts and its esters are dangerous to man and the environment, and in particular to the aquatic environment. They have been classified by the EC as category 3 carcinogens (possibly carcinogenic to humans)

<b>Belize</b>	
<b>Control Action</b> <b>Effective</b> <b>Uses still allowed</b> <b>Reasons for control action</b>	Severely restricted 1985 Wood preservation purposes only by approved and certified establishments and personnel Oncogen, mutagen, teratogen, high dermal toxicity

<b>China</b>	
<b>Control Action</b>  <b>Effective</b>	Pentachlorophenol has been banned for registration and production, sale and use as pesticide. Uses are only allowed as wood preservative. 1982

<sup>1</sup> Members of the European Union (EU): Austria, Belgium, Denmark, Finland, France, Germany, Greece, Ireland, Italy, Luxembourg, Netherlands, Portugal, Spain, Sweden, United Kingdom  
Members of the European Economic Agreement (EEA): Iceland, Liechtenstein, Norway

<b>Uses still allowed</b>	Use is still allowed as germicide on woods. However, areas and methods of approved application as stated in "Bulletin of Pesticide Registration" should be observed.
<b>Reasons for control action</b>	These measures were taken because pentachlorophenol is a highly toxic germicide and its abuse is severely harmful to human health
<b>Date of notification</b>	

## ANNEX 2

### Alternatives

*The following alternatives were noted by countries reporting import decisions under the PIC procedure:*

Country	
<b>Austria</b>	Many alternatives for designated purposes
<b>India</b>	Paraquat as herbicide. TCMTB as fungicide.
<b>Indonesia</b>	Many less toxic wood preservatives already registered

*It is essential that before a country considers substituting any of these reported alternatives, it ensures that the use is relevant to their national needs. A first step may be to contact the DNA in the country where the alternative has been reported (see address: Annex 3). It will then be necessary to determine the compatibility with national crop protection practices.*

## ANNEX 3

<b>List of Pesticide DNAs in Countries Reporting Control Actions or Alternatives</b>
--

<b>Belgium</b>	<b>CP</b>	Service Maîtrise des risques Section Pesticides (bureau 2/309) Ministère de la santé publique et de l'environnement Cité Administrative de l'Etat 1010 Bruxelles (Attn. Mr. R. Huysman)	Tel: 32 2 2104881 Fax: 32 2 2104884
<b>Belize</b>	<b>P</b>	The Secretary Pesticides Control Board Department of Agriculture Central Farm Cayo	Tel: 501-92-2640 Fax: 501-92-2640 Tlx: 102 Foreign Bz
	<b>C</b>	Mr. Carlos Guerra Sanitation Engineer Ministry of Public Health Public Health Bureau Belize City	
<b>China</b>	<b>P</b>	The Director Institute for the Control of Agrochemicals (ICAMA) Ministry of Agriculture Liang Ma Qiao, Chaoyang Beijing 100026	Tel/Fax: 86 010 5025929
	<b>CP</b>	National Environmental Protection Agency (NEPA) No. 115, Xizhimennei Nanxiaojie Beijing 100035 (Attn.: Mrs Sun Lijin)	Tel: 8329911, Ext. 3555/3609 Tlx: 222359 NEPA CN Fax: 8328013
<b>India</b>	<b>P</b>	The Director/Deputy Secretary Plant Protection Division Dept. of Agriculture & Co-op. Room No. 244-A Krishi Bhavan, New Delhi	
	<b>C</b>	Adviser (Chemicals) Dept. of Chemicals & Petrochemicals Ministry of Chemicals & Fertilizers Shastri Bhavan Rajendra Prasad Road New Delhi - 110001	Tel: 91 (11) 385736/382575 Tlx: 62455 Fax: 91 (11) 382604/337223
<b>Indonesia</b>	<b>P</b>	Chairman Pesticides Committee Direktorat Bina Perlindungan Tanaman Jln. AUP. Pasar Minggu Jakarta Selatan	Tel: 62 (21) 7805652/7806213 Fax: 62 (21) 7805652
	<b>CP</b>	Ms. Masnellyarti Hilman Bapedal Offices Arthaloa Bldg., 11th Floor Jl. Jend. Sudirman No. 2 Jakarta Pusat	Tel: (021) 583918 Tlx: 62 21 583918 Fax: (021) 5703365

<b>Kuwait</b>	<b>P</b>	The Director Plant Wealth Department The Public Authority for Agriculture Affairs & Fish Res. P.O. Box 21422 13075 Safat	Tel: (965) 2452790, 2456835/36 Tlx: 46408 EP CNCL KT Fax: (965) 2421993-2456836
<b>New Zealand</b>	<b>CP</b>	Mr. D.W. Lunn Chief Scientist (Pesticides) Agricultural Compounds Unit Ministry of Agriculture & Fisheries P.O. Box 40-063 Upper Hutt	Tel: 064 4 528-6089 Fax: 064 4 528-4675
<b>Sweden</b>	<b>CP</b>	National Chemicals Directorate Attn.: Mr. Ule Johansson P.O. Box 1384 171 27 Solna	Tel: 46 (8) 730 6004 Tlx: 10460 AMS S Fax: 46 (8) 735 7698
<b>Switzerland</b>	<b>CP</b>	Service des affaires internationales Office fédéral de l'environnement, des forêts et du paysage (OFEFP) Hallwylstr. 4, 3003 Berne	Tel: 41 31 322 99 73 Fax: 41 31 322 99 81 Tlx: 91 23 04
<b>USA</b>	<b>CP</b>	The Assistant Administrator for Pesticides and Toxic Substances Environmental Protection Agency 401 M St. S.W. Washington DC 20460	Tel: 1 202 260 2902 Fax: 1 202 260 1847 Tlx: 892758 EPA WSH

	<b>C</b>	Industrial and consumer product chemicals	
	<b>P</b>	Pesticides	
	<b>CP</b>	Pesticides, industrial and consumer product chemicals	

